

Iron Deposits after Orthokeratology

Iron Deposits after Orthokeratology

Dr. Andreas Berke
Peter Bruckmann
School of Optometry
Cologne, Germany

GLOBAL
SYMPOSIUM

Purpose

- To study the frequency of iron deposits after OK.
- To study the impact of iron deposits on vision.

GLOBAL
SYMPOSIUM

Patients

- 32 patients, 64 eyes
- Age 29.2 ± 5.7 years
- Mean pre-OK myopia: 3.25 ± 1.25 D
- Orthokeratology for 14 ± 5 months
- OK lenses: Technolens (Switzerland), ProCornea/MPGE (Netherlands, Germany)

GLOBAL
SYMPOSIUM

Dr. Andreas Berke
Cologne, Germany

Iron Deposits after Orthokeratology


Methods

- Slit lamp examination (Zeiss SL 160)
- Visual acuity (Standard tests)
- Contrast sensitivity (Freiburger Visus Test)

Results


- 16 of 64 eyes (25 %) with arcuate iron deposits in the lower cornea (reverse zone of OK lens).
- First appearance 2 months after start of OK
- Mean: 5 ± 2 months
- No progression after first appearance

Spherical Equivalents before OK




Iron Deposits after Orthokeratology


Changes of Corneal Radius



Ferritin Line after OK



Ferritin Line after OK




Iron Deposits after Orthokeratology

Iron Deposits or Ferritin Lines


- Ferritin lines occur seldom in isolation. They are associated with an underlying corneal problem.
 - *Hudson-Stähli's line (physiological in elderly)*
 - *Ferry's line (anterior to a filtering bleb)*
 - *Stocker's line (anterior to a pterygium)*
 - *Fleischer's ring: Keratoconus*
 - *Foreign body*
 - *Other (Diseases that distort the surface of the cornea, Keratoplasty, Intrastromal Corneal Ring, LASIK)*

GLOBAL
ORTHOKERATOLOGY

Ferritin Line



LASIK



Keratoconus

GLOBAL
ORTHOKERATOLOGY


Ferritin line after OK

- Iron deposits are related to poor tear film exchange
 - *Ferritin line appears within the reverse zone of OK lenses.*
 - *Stagnant tear film in the reverse zone*
- OK results in distortion of corneal surface
 - *Iron uptake may be facilitated*
- No progression after a few months because of stabilization of the corneal epithelium.

GLOBAL
ORTHOKERATOLOGY

Iron Deposits after Orthokeratology

Iron Storage: Ferritin




- Iron storage within ferritin
 - *Ferritin is composed of 24 identical protein subunits that form a hollow shell.*
 - *4500 iron ions inside where they form small crystallites along with phosphate and hydroxide ions*

GLOBAL
ORTHOKERATOLOGY

Ferritin

- Suppression of Fenton reaction:



- Storage of ferric form: Fe_2O_3 , FePO_4 and $\text{Fe}(\text{OH})_3$
- Protects nucleus of epithelial cells from UV damage.

GLOBAL
ORTHOKERATOLOGY

Visual functions after one year

- Iron deposits do not interfere with vision.
 - *Visual acuity: 20/20 and better*
 - *Contrast sensitivity: 45*
- No loss of visual acuity and contrast sensitivity compared to pre-OK values

GLOBAL
ORTHOKERATOLOGY

Iron Deposits after Orthokeratology

Conclusions

- Iron deposits occur frequently after a few months of OK.
- There is no progression.
- The degree of corneal reshaping seems to influence iron deposits.
- Iron deposits do not interfere with vision.
- They seem to be harmless.

GLOBAL
SYMPOSIUM

Iron Deposits after Orthokeratology

Dr. Andreas Berke
Peter Bruckmann
School of Optometry
Cologne, Germany

GLOBAL
SYMPOSIUM
